

УДК 514.764.21, 514.765, 512.813

DOI 10.46698/i4125-5722-6924-j

ON EXAMPLES OF GEODESIC ORBIT PSEUDO-RIEMANNIAN MANIFOLDS

I. G. Markina¹, Yu. G. Nikonorov² and K. Furutani³

¹ Department of Mathematics, University of Bergen,
P. O. Box 7803, N-5020 Bergen, 41 Allégaten, Norway;

² Southern Mathematical Institute of VSC RAS,
53 Vatutina St., Vladikavkaz 362025, Russia;

³ Osaka Central Advanced Mathematical Institute, Osaka Metropolitan University,
Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan

E-mail: irina.markina@uib.no, nikonorov2006@mail.ru, kf46089@gmail.com

Dedicated to the memory of Professor Semën Samsonovich Kutateladze

Abstract. A pseudo-Riemannian manifold (M, g) is called a geodesic orbit manifold if any geodesic γ of M is an orbit of a 1-parameter subgroup of the full isometry group of (M, g) . This terminology in the case of Riemannian manifolds was introduced in 1991 by O. Kowalski and L. Vanhecke, who initiated a systematic study of spaces $(M = G/H, g)$, where G is an isometry group and H is an isotropy subgroup. It should be noted that symmetric spaces, weakly symmetric spaces, naturally reductive homogeneous spaces, normal homogeneous spaces, generalized normal homogeneous spaces (but not only) are subclasses of geodesic orbit pseudo-Riemannian spaces. In this paper, we present examples of geodesic orbit pseudo-Riemannian manifolds. The examples are special 15-dimensional pseudo H -type Lie groups, i. e., 2-step nilpotent Lie groups of Heisenberg type equipped with a left invariant pseudo-Riemannian metric. To construct the corresponding examples, results on the structure of the Lie groups under consideration were used.

Keywords: geodesic orbit Riemannian manifolds, geodesic orbit pseudo-Riemannian manifolds, H -type Lie groups.

AMS Subject Classification: 53C50, 53C30, 22E25.

For citation: Markina, I. G., Nikonorov, Yu. G. and Furutani, K. On Examples of Geodesic Orbit Pseudo-Riemannian Manifolds, *Vladikavkaz Math. J.*, 2026, vol. 28, no. 1, pp. 108–121. DOI: 10.46698/i4125-5722-6924-j.

1. Introduction

A pseudo-Riemannian manifold (M, g) is called a geodesic orbit manifold if any geodesic γ of M is an orbit of a 1-parameter subgroup of the full isometry group of (M, g) . This terminology in the case of Riemannian manifolds was introduced in [1] by O. Kowalski and L. Vanhecke, who initiated a systematic study of spaces $(M = G/H, g)$, where G is an isometry group and H is an isotropy subgroup. We refer to [2–6] for expositions on general properties of geodesic orbit Riemannian manifolds and historical surveys. Some important results on geodesic orbit pseudo-Riemannian spaces were obtained in [7–11]. It should be noted that symmetric spaces, weakly symmetric spaces, naturally reductive homogeneous spaces, normal homogeneous spaces, generalized normal homogeneous spaces (but not only) are subclasses of geodesic orbit pseudo-Riemannian spaces.

The pseudo-Riemannian H -type nilmanifold is a 2-step nilpotent H -type Lie group endowed with a left invariant pseudo-Riemannian metric. The Lie algebra \mathfrak{n} of a pseudo-Riemannian H -type nilmanifold carries a non-degenerate scalar product $\langle \cdot, \cdot \rangle$, whose restriction $\langle \cdot, \cdot \rangle_{\mathfrak{z}}$ on the centre \mathfrak{z} of the Lie algebra \mathfrak{n} is also non-degenerate. The orthogonal complement \mathfrak{v} to the center is a representation space for the Clifford algebra $\text{Cl}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$, such that the representation map is the dual to the adjoint map on the Lie algebra, see Section 2.

The problem of classification of geodesic orbit and naturally reductive pseudo-Riemannian nilmanifolds of H -type is very relevant. The results on Riemannian H -type nilmanifolds (the metric is positive definite) were obtained by A. Kaplan in [12] and C. Riehm in [13] (see Theorem 1) and the results on pseudo-Riemannian H -type nilmanifolds, when the space \mathfrak{v} is a minimal dimensional admissible Clifford module was obtained in [14].

Theorem 1 [12, 13]. *Let N be H -type Lie group (supplied with a left invariant Riemannian metric) with the H -type algebra $\mathfrak{n} = \mathfrak{z} \oplus \mathfrak{v}$, $m = \dim(\mathfrak{z})$, $n = \dim(\mathfrak{v})$, and the center \mathfrak{z} . Then N is geodesic orbit if and only if one of the following three conditions holds:*

- 1) $m = 1, 2, 3$ and n is any possible;
- 2) $m = 5, 6$ and $n = 8$;
- 3) $m = 7, n = 8, 16, 24$ and \mathfrak{v} is an isotypic Clifford module (in this case it is equivalent to the following property: if Z_1, Z_2, \dots, Z_7 is an orthonormal basis of \mathfrak{z} , the linear transformation $X \mapsto J_{Z_1} J_{Z_2} \cdots J_{Z_7}(X)$ of \mathfrak{v} is either Id or $-\text{Id}$).

Moreover, N is naturally reductive if and only if $m = 1$ or $m = 3$.

Recently, an analogue of this theorem was obtained for pseudo-Riemannian metrics.

Theorem 2 [14]. *Let $N_{r,s}$ be an H -type Lie group, where (r, s) , $s \geq 1$, is the signature of the left invariants pseudo-Riemannian metric restricted to the center of the group. Let $\mathfrak{n}_{r,s} = \mathfrak{z} \oplus \mathfrak{v}$ be the Lie algebra of $N_{r,s}$, where \mathfrak{z} is the centre and \mathfrak{v} is a minimal admissible module for the Clifford algebra $\text{Cl}(\mathbb{R}^{r,s})$ orthogonal to \mathfrak{z} . Then the following four assertions hold:*

- 1) $N_{r,s}$ is naturally reductive (hence, geodesic orbit) if and only if $(r, s) \in \{(0, 1), (1, 2)\}$;
- 2) If $N_{r,s}$ is geodesic orbit but not naturally reductive, then $(r, s) = (3, 4)$;
- 3) $N_{3,4}$ is a geodesic orbit pseudo-Riemannian manifold;
- 4) $N_{r,s}$ with $(r, s) \notin \{(0, 1), (1, 2), (3, 4)\}$ is not geodesic orbit pseudo-Riemannian manifold.

It is important that item 3) of the above theorem is proved in [14] using a system of symbolic computations that allows one to calculate and evaluate the ranks of a number of auxiliary matrices to prove that each geodesic can be represented as an orbit of a 1-parameter isometry group of the manifold under study. This kind of reasoning proves the existence of the required one-parameter group, but does not always allow one to specify it explicitly.

The main goal of this note is to provide an alternative proof of the corresponding result, which also provides explicit expressions for the corresponding one-parameter isometry groups, see Theorem 4.

The paper is organized as follows. In Section 2, we recall some important results on pseudo-Riemannian geodesic orbit metrics on nilpotent Lie groups. The main role here is played by the notion of the transitive normalizer condition, which Riemannian version was used by C. Gordon in order to study geodesic orbit Riemannian metric on nilpotent Lie groups. We also recall some important properties of pseudo H -type Lie groups, their isometry and automorphism groups. In Section 3, we show the geodesic orbit property for the 15-dimensional pseudo-Riemannian H -type nilmanifold $N_{3,4}(\mathfrak{v}_{\min})$. Section 4 is dedicated to the same property for the Riemannian H -type nilmanifold $N_{7,0}(\mathfrak{v}_{\min})$.

2. Definitions and Notation

2.1. Geodesic orbit pseudo-Riemannian manifolds.

DEFINITION 1. A pseudo-Riemannian homogeneous reductive manifold $(G/H, g)$ is called *geodesic orbit* if any geodesic through the point eH is an orbit of some 1-parameter isometry group of $(G/H, g)$.

Let (N, g) be a 2-step pseudo-Riemannian nilpotent Lie group with the Lie algebra \mathfrak{n} and $\langle \cdot, \cdot \rangle$ the scalar product (a symmetric non-degenerate bilinear form) generating the pseudo-Riemannian left invariant metric g . We call such groups *pseudo-Riemannian nilmanifolds*. If the centre \mathfrak{z} of \mathfrak{n} is non-degenerate with respect to $\langle \cdot, \cdot \rangle$, then we write $\mathfrak{n} = \mathfrak{z} \oplus \mathfrak{v}$, where $\mathfrak{v} = \mathfrak{z}^\perp$ relative to $\langle \cdot, \cdot \rangle$. In this case, \mathfrak{v} is also non-degenerate, see [15, Lemma 2.60].

Whenever (N, g) is connected simply connected, we do not distinguish between the group of automorphisms of the nilmanifold (N, g) and of its Lie algebra $\mathfrak{n} = \mathfrak{z} \oplus \mathfrak{v}$. Note that each skew-symmetric derivation of \mathfrak{n} leaves each of \mathfrak{v} and \mathfrak{z} invariant. For any $Z \in \mathfrak{z}$, we consider the operator

$$J_Z: \mathfrak{v} \rightarrow \mathfrak{v}, \quad \text{such that} \quad \langle J_Z(X), Y \rangle = \langle [X, Y], Z \rangle, \quad X, Y \in \mathfrak{v}. \quad (1)$$

The map J_Z is skew-symmetric and $J_Z(Y) = (\text{ad } Y)'(Z)$, where $(\text{ad } Y)'$ is dual to $\text{ad } Y$ with respect to $\langle \cdot, \cdot \rangle$. The map $J: \mathfrak{z} \rightarrow \mathfrak{so}(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}})$, sending $Z \mapsto J_Z$ is linear. The group of isometries of the nilmanifold (N, g) is given by

$$H = \{(\varphi, \psi) \in O(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}}) \times O(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}}) : \psi J_Z \psi^{-1} = J_{\varphi(Z)}, Z \in \mathfrak{z}\}, \quad (2)$$

while its Lie algebra is

$$\begin{aligned} \mathfrak{h} &= \text{Der}(\mathfrak{n}) \cap \mathfrak{so}(\mathfrak{n}, \langle \cdot, \cdot \rangle) \\ &= \{D = (C, A) \in \mathfrak{so}(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}}) \times \mathfrak{so}(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}}) : [A, J_Z] = J_{C(Z)}, Z \in \mathfrak{z}\}. \end{aligned} \quad (3)$$

The next result is well known, see e. g. Corollary 3.5 in [16].

Proposition 1. *Let (N, g) be a pseudo-Riemannian nilmanifold with non-degenerate center. Then the connected isometry group of (N, g) is $N \rtimes H$, where N is the set of left translations by elements of N and the isotropy subgroup H is given by the isometric automorphisms (2) with Lie algebra \mathfrak{h} as in (3).*

In this case, the isotropy group of (N, g) at the identity element e is exactly H with the embedding $a \in H \mapsto (e, a) \in N \rtimes H$. The following Proposition summarizes well known Geodesic Lemma [10], its reformulation in the spirit of [17], and consideration done in [14] for the setting of H -type nilmanifolds, see definition in Section 2.2.

Proposition 2. *A pseudo-Riemannian H -type nilmanifold (N, g) is geodesic orbit if for any $z_0 \in \mathfrak{z}$ and $x_0 \in \mathfrak{v}$ there is $D = (C, A) \in \mathfrak{h}$, such that*

$$A(x_0) = J_{z_0}(x_0), \quad C(z_0) = 0, \quad [A, J_w] = J_{C(w)} \quad \text{for all } w \in \mathfrak{z}. \quad (4)$$

A special case of geodesic orbit pseudo-Riemannian spaces are naturally reductive homogeneous pseudo-Riemannian spaces. We refer to [14] and references therein for more detailed study.

DEFINITION 2. Let $(M = G/H, g)$ be a homogeneous reductive pseudo-Riemannian manifold. Then M is said to be a naturally reductive if there is a reductive decomposition $\mathfrak{g} = \mathfrak{h} \oplus \mathfrak{m}$, such that

$$\langle [T, Q]_{\mathfrak{m}}, R \rangle = \langle Q, [T, R]_{\mathfrak{m}} \rangle \quad (5)$$

for the corresponding scalar product and any $T, Q, R \in \mathfrak{m}$.

Theorem 3 [18, Theorem 3.2]. *Let (N, g) denote a 2-step pseudo-Riemannian nilmanifold with a non-degenerate center. Assume that the map $J: \mathfrak{z} \rightarrow \mathfrak{so}(\mathfrak{v})$ is injective, see (1). Then the metric is naturally reductive with respect to $G = N \rtimes H$ (see Proposition 1), if and only if*

- (i) $\mathbf{V} = J(\mathfrak{z})$ is a Lie subalgebra of $\mathfrak{so}(\mathfrak{v})$ and
- (ii) $[J(Z_1), J(Z_2)] = J(\tau_{Z_1}(Z_2))$, where $\tau_{Z_1} \in \mathfrak{so}(\mathfrak{z})$ for any $Z_1 \in \mathfrak{z}$.

Since the map J is supposed to be injective, the map τ can be easily recovered from (ii).

REMARK 1. Using Theorem 3, it is easy to prove that the pseudo-Riemannian H -type nilmanifold $N_{3,4}(\mathfrak{v}_{\min})$, as well as the corresponding Riemannian H -type nilmanifold $N_{7,0}(\mathfrak{v}_{\min})$, that are studied further in this note, are not naturally reductive.

2.2. Pseudo H -type Lie groups. Now, we recall some important properties of pseudo H -type Lie groups. Let (N, g) be a 2-step pseudo-Riemannian nilmanifold and $\mathfrak{n} = \mathfrak{z} \oplus \mathfrak{v}$ be its Lie algebra endowed with a scalar product $\langle \cdot, \cdot \rangle$ making the center non-degenerate. We identify $(\mathfrak{z}, \langle \cdot, \cdot \rangle_{\mathfrak{z}})$ with the pseudo Euclidean vector space $\mathbb{R}^{r,s} = (\mathbb{R}^{r+s}, \langle \cdot, \cdot \rangle_{r,s})$, where

$$\langle z, w \rangle_{r,s} = \sum_{i=1}^r z_i w_i - \sum_{j=1}^s z_{r+j} w_{r+j}, \quad z, w \in \mathbb{R}^{r+s}.$$

If the linear operator $J: \mathfrak{z} \rightarrow \mathfrak{so}(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}})$ is defined by for any $z \in \mathfrak{z}$ by (1), and satisfies $J_z^2(x) = -\langle z, z \rangle_{r,s} x$ for any $z \in \mathbb{R}^{r,s}$ and all $x \in \mathfrak{v}$, then $\mathfrak{n} = \mathfrak{n}_{r,s}$ is called the pseudo H (eisenberg)-type Lie algebra. We denote by $N_{r,s}$ the connected simply connected Lie group, whose Lie algebra is the pseudo H -type Lie algebra $\mathfrak{n}_{r,s}$. The H -type Lie algebras $N_{r,0}$ with a positive definite scalar product were introduced in [19] and with an arbitrary indefinite scalar product in [20], see also [21].

These Lie algebras are related to the representations of the Clifford algebras in the following way. Let $J: \text{Cl}(\mathbb{R}^{r,s}) \rightarrow \text{End}(\mathfrak{v})$ be a representation of the Clifford algebra $\text{Cl}(\mathbb{R}^{r,s})$ generated by the pseudo Euclidean vector space $\mathbb{R}^{r,s}$. If there is a scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ on the representation space \mathfrak{v} (Clifford module) such that the linear map J_z is skew-symmetric for any $z \in \mathbb{R}^{r,s}$; that is

$$\langle J_z(x), y \rangle_{\mathfrak{v}} = -\langle x, J_z(y) \rangle_{\mathfrak{v}}, \quad z \in \mathbb{R}^{r,s}, \quad x, y \in \mathfrak{v},$$

then we get a pseudo H -type Lie algebra with the commutators defined in (1), see details in [20, 22, 23]. The scalar product $\langle \cdot, \cdot \rangle_{\mathfrak{v}}$ in this case is called admissible and \mathfrak{v} is called admissible (Clifford) module.

It is important that, see e.g. [20], or [23, Proposition 2.2.2], the signature of the scalar product space $(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}})$ is neutral and $(\mathfrak{v}, \langle \cdot, \cdot \rangle_{\mathfrak{v}})$ is isometric to $\mathbb{R}^{l,l}$ for some $l \in \mathbb{N}$, if $s > 0$, whereas the corresponding signature is either $(l, 0)$ or $(0, l)$ for some $l \in 2\mathbb{N}$, if $s = 0$.

3. Pseudo-Riemannian H -Type Nilmanifold $N_{3,4} = N_{3,4}(\mathfrak{v}_{\min})$

Although the fact that a pseudo-Riemannian nilmanifold of H -type $N_{3,4}(\mathfrak{v}_{\min})$, where the complement of the centre is the minimal admissible modulus, is a geodesic orbit manifold is proved in [14], we give here an alternative proof of the following theorem.

Theorem 4. *The pseudo-Riemannian H -type nilmanifold $N_{3,4}(\mathfrak{v}_{\min})$ is a geodesic orbit manifold. Each geodesic is realized as an orbit of a one-parameter subgroup which initial conditions form an eight dimensional affine space.*

It is known that the Clifford algebra $\text{Cl}(\mathbb{R}^{3,4})$ has two non-equivalent minimal admissible 8-dimensional modules \mathfrak{v}_{\min}^+ and \mathfrak{v}_{\min}^- . Moreover, the corresponding 15-dimensional pseudo

H -type Lie algebras $N_{3,4}(\mathfrak{v}_{\min}^+)$ and $N_{3,4}(\mathfrak{v}_{\min}^-)$ are isomorphic and isometric, see [24, Theorem 12]. Therefore, it suffices to check only one pseudo H -type nilmanifold $N_{3,4}(\mathfrak{v}_{\min}^-) = N_{3,4}(\mathfrak{v}_{\min})$. The left invariant metric on $N_{3,4}(\mathfrak{v}_{\min})$ is generated by the scalar product $\langle \cdot, \cdot \rangle_{3,4} + \langle \cdot, \cdot \rangle_{4,4}$ on the Lie algebra $\mathfrak{n}_{3,4} \cong \mathbb{R}^{3,4} \oplus \mathbb{R}^{4,4}$. We write $(z_0, x_0) \in \mathfrak{z} \oplus \mathfrak{v}_{\min} \cong \mathbb{R}^{3,4} \oplus \mathbb{R}^{4,4}$ for an initial vector for a geodesic. For different type of the initial vectors we will use distinguished bases for the horizontal space \mathfrak{v}_{\min} and for the centre \mathfrak{z} . We start from a basis for the centre.

3.1. Central component z_0 of initial conditions.

(C1) If $\|z_0\|^2 = \langle z_0, z_0 \rangle_{\mathbb{R}^{3,4}} = 1$, then we set

$$\{Z_1 = z_0, Z_2, \dots, Z_7\},$$

as an orthonormal basis, satisfying

$$\|Z_1\|^2 = \|Z_2\|^2 = \|Z_3\|^2 = 1, \quad \|Z_4\|^2 = \|Z_5\|^2 = \|Z_6\|^2 = \|Z_7\|^2 = -1. \quad (6)$$

(C2) If $\|z_0\|^2 = \langle z_0, z_0 \rangle_{\mathbb{R}^{3,4}} = -1$, then we set

$$\{Z_1, Z_2, Z_3, Z_4 = z_0, Z_5, Z_6, \dots, Z_7\},$$

such that (6) holds.

(C3) If $\|z_0\|^2 = \langle z_0, z_0 \rangle_{\mathbb{R}^{3,4}} = 0$, then first we fix an arbitrary orthonormal basis $\{z_1^*, \dots, z_7^*\}$, satisfying (6) and write

$$z_0 = \sum_{i=1}^3 \theta_i z_i^* + \sum_{i=4}^7 \theta_i z_i^* := z_+^* + z_-^*.$$

Then, without loss of generality, we can assume $\sum_{i=1}^3 \theta_i^2 = \sum_{i=4}^7 \theta_i^2 = 1$. Since $\langle z_+^*, z_-^* \rangle_{\mathbb{R}^{3,4}} = 0$ we can select an orthonormal basis

$$\{Z_1 = z_+^*, Z_2, Z_3, Z_4 = z_-^*, Z_5, Z_6, Z_7\},$$

still satisfying (6). With these basis vectors we can denote $z_0 = Z_1 + Z_4$.

In what follows we use the construction of basis on the centre as above.

REMARK 2. In any of the above constructions, the operators J_{Z_j} , $j = 1, \dots, 7$, span a 7-dimensional subspace in $J_{\mathfrak{z}} \subset \mathfrak{so}(4, 4)$, nevertheless the vector space $\mathbf{V} = J_{\mathfrak{z}}$ is not a Lie subalgebra of $\mathfrak{so}(4, 4)$. Hence, $N_{3,4}(\mathfrak{v}_{\min})$ is not a naturally reductive manifold, see Theorem 3.

3.2. Horizontal component x_0 of initial conditions and reduction by conjugate Clifford action. Fix $z_0 \in \mathfrak{z}$ and choose a suitable basis as was described in Section 3.1. Now we have three cases:

$$\textbf{(H1)} \quad \|x_0\|^2 = \langle x_0, x_0 \rangle_{\mathbb{R}^{4,4}} = 1, \quad \textbf{(H2)} \quad \|x_0\|^2 = -1, \quad \textbf{(H3)} \quad \|x_0\|^2 = 0.$$

Before we consider these three options we let $v \in \mathfrak{v}_{\min}$ be such that $\|v\|^2 = 1$ and

$$J_{Z_1 Z_2 Z_4 Z_5}(v) = v, \quad J_{Z_1 Z_2 Z_6 Z_7}(v) = v, \quad J_{Z_1 Z_3 Z_5 Z_7}(v) = v, \quad J_{Z_1 Z_2 Z_3}(v) = v. \quad (7)$$

The condition $J_{Z_1 Z_2 Z_3}(v) = v$ is equivalent to $J_{Z_1 Z_2 Z_3 Z_4 Z_5 Z_6 Z_7}(v) = -v$. This choice fixes the module $\mathfrak{v}_{\min} = \mathfrak{v}_{\min}^-$. Moreover, the choice of v as in (7) allows us to use the following orthonormal basis for the module \mathfrak{v}_{\min} :

$$\begin{aligned} X_1 = v, \quad X_2 = J_{Z_1}(v), \quad X_3 = J_{Z_2}(v), \quad X_4 = J_{Z_3}(v), \\ X_5 = J_{Z_4}(v), \quad X_6 = J_{Z_5}(v), \quad X_7 = J_{Z_6}(v), \quad X_8 = J_{Z_7}(v). \end{aligned} \quad (8)$$

REMARK 3. Once we fix an orthonormal basis $\{z_i\}$ in $\mathbb{R}^{3,4}$ (or in general in $\mathbb{R}^{r,s}$), the number of a set of generating positive involutions $\{p_i\}$ of a maximal abelian subgroup S consisting of positive involutions, which is a subgroup in the finite group $\mathbb{G} = \mathbb{G}(\{z_i\})$ generated by $\{z_i\}$, is determined by the pair (r, s) . In the case $(r, s) = (3, 4)$ it is 4, that is one set of them is listed in (7). It should be noted that in our case the correspondence $z \mapsto J_z$ is not one-to-one. This is common in all the cases of the Clifford algebra being not irreducible. In such case we have two irreducible modules and they are distinguished by the action of the volume form $J_{z_1} \cdots J_{z_{r+s}}$. As we can see easily that in our case $J_{z_1} \cdots J_{z_7}$ is in the center of the group \mathbb{G} , so that $J_{z_1} J_{z_2} J_{z_3}(v) = v$ is not only equivalent to $J_{z_1} \cdots J_{z_7}(v) = -v$, it is also equivalent to $J_{z_1} \cdots J_{z_7} \equiv -\text{Id}$ as an operator acting on the space generated by $\{J_{z_i}(v)\}$. So we may distinguish two irreducible modules \mathfrak{v}_{\min}^\pm by the action of the volume form $J_{z_1} \cdots J_{z_7} = \pm \text{Id}$, see details in [24].

Let $x_0 \in \mathfrak{v}_{\min}$ be as in **(H1)** above. Let $T \in SO(4, 4)$ be such that $x_0 = T(v)$ and consider the conjugate Clifford action $\tilde{J}_w := T^{-1} J_w T = T^\tau J_w T$, where T^τ is the transpose to T with respect to the scalar product $\langle \cdot, \cdot \rangle_{4,4}$. Then, a geodesic determined by the initial condition (z_0, x_0) is homogeneous if there is a pair $D = (C, A)$ such that

$$A(x_0) = J_{z_0}(x_0), \quad C(z_0) = 0, \quad [A, J_w] = J_{C(w)}, \quad \text{for all } w \in \mathbb{R}^{3,4}. \quad (9)$$

Lemma 1. *The condition (9) is equivalent to*

$$\tilde{A}(v) = T^{-1} A T(v) = T^{-1} J_{z_0} T(v) = \tilde{J}_{z_0}(v), \quad C(z_0) = 0, \quad [\tilde{A}, \tilde{J}_w] = \tilde{J}_{C(w)}, \quad (10)$$

for all $w \in \mathfrak{z} \cong \mathbb{R}^{3,4}$.

◁ In fact, from (9)

$$\begin{aligned} T^{-1} A T(v) &= T^{-1} A(x_0) = T^{-1} J_{z_0}(x_0) = T^{-1} T \tilde{J}_{z_0} T^{-1}(x_0) \\ &= \tilde{J}_{z_0}(v), [T^{-1} A T, T^{-1} J_w T] = T^{-1} [A, J_w] T = T^{-1} J_{C(w)} T = \tilde{J}_{C(w)}, \end{aligned}$$

and vice versa. ▷

We denote the H -type group $N_{3,4}(\mathfrak{v}_{\min})$ related to the Clifford action $J_z: \mathfrak{v}_{\min} \rightarrow \mathfrak{v}_{\min}$ and the H -type group $N_{3,4}^T(\mathfrak{v}_{\min})$ related to the Clifford action $\tilde{J}_z: \mathfrak{v}_{\min} \rightarrow \mathfrak{v}_{\min}$, $\tilde{J}_z = T^{-1} J_z T$. Then the Lie bracket $[x, y]^T$ defined by \tilde{J}_z is given by

$$\langle z, [x, y]^T \rangle_{\mathbb{R}^{3,4}} = \left\langle \tilde{J}_z(x), y \right\rangle_{\mathfrak{v}_{\min}}.$$

Lemma 2. *The transformation $T \times \text{Id}: N_{3,4}(\mathfrak{v}_{\min}) \rightarrow N_{3,4}^T(\mathfrak{v}_{\min})$ is an isometric isomorphism.*

◁ We can see this by the equality:

$$\begin{aligned} \langle z, [T(x), T(y)] \rangle_{\mathfrak{v}_{\min}} &= \langle J_z(T(x)), T(y) \rangle_{\mathfrak{v}_{\min}} \\ &= \langle T^\tau J_z T(x), y \rangle_{\mathfrak{v}_{\min}} = \left\langle \tilde{J}_z(x), y \right\rangle_{\mathfrak{v}_{\min}} = \langle z, [x, y]^T \rangle_{\mathbb{R}^{3,4}}, \end{aligned}$$

that proves the lemma. ▷

Let $x_0 \in \mathfrak{v}_{\min}$ be as in **(H2)** and $v \in \mathfrak{v}_{\min}$ as in (7). We find $T \in SO(4, 4)$, $T: \mathfrak{v}_{\min} \rightarrow \mathfrak{v}_{\min}$, such that $T(x_0) = J_{Z_4}(v)$. Then condition (9) is equivalent to

$$A(J_{Z_4}(v)) = J_{z_0}(v), \quad C(z_0) = 0, \quad [A, J_w] = J_{C(w)} \quad \text{for all } w \in \mathbb{R}^{3,4}.$$

by changing the Clifford action from J_z to $\tilde{J}_z = T^{-1}J_zT$.

Let $x_0 \in \mathfrak{v}_{\min}$ be as in **(H3)** and $v \in \mathfrak{v}_{\min}$ as in (7). We fix an orthonormal basis $\{Z_i\}_{i=1}^7$ as was described in **(C3)**, and express x_0 in the basis (8) as follows

$$x_0 = \sum_{i=0}^3 \theta_i J_{Z_i}(v) + \sum_{i=4}^7 \theta_i J_{Z_i}(v) = x_0^+ + x_0^-, \quad \sum_{i=0}^3 \theta_i^2 = \sum_{i=4}^7 \theta_i^2 = 1.$$

Since $\langle x_0^+, x_0^- \rangle_{\mathfrak{v}_{\min}} = 0$, we can find an orthogonal transformation $T \in SO(4, 0) \times SO(0, 4) \subset SO(4, 4)$, such that $T(x_0^+) = v$ and $T(x_0^-) = J_{Z_4}(v)$. Then for $\tilde{J}_z = T^{-1}J_zT$ we obtain

$$\begin{aligned} T^{-1}AT(v + J_{Z_4}(v)) &= T^{-1}J_{z_0}T(v + J_{z_4}(v)), \quad C(z_0) = 0, \\ T^{-1}[A, J_w]T &= [T^{-1}AT, T^{-1}J_wT] = T^{-1}J_{C(w)}T \quad \text{for any } w \in \mathbb{R}^{3,4}, \end{aligned} \quad (11)$$

which is equivalent to condition (9).

The discussions above can be summarized in the statement that an initial vector (z_0, x_0) for a geodesic can be restricted to the following nine cases.

$$\left. \begin{array}{l} Z_1 \oplus v, \\ Z_4 \oplus v, \\ (Z_1 + Z_4) \oplus v \end{array} \right\} \in \mathbb{R}^{3,4} \oplus \mathfrak{v}_{\min}, \quad (12)$$

$$\left. \begin{array}{l} Z_1 \oplus J_{Z_4}(v), \\ Z_4 \oplus J_{Z_4}(v), \\ (Z_1 + Z_4) \oplus J_{Z_4}(v) \end{array} \right\} \in \mathbb{R}^{3,4} \oplus \mathfrak{v}_{\min}, \quad (13)$$

$$\left. \begin{array}{l} Z_1 \oplus (v + J_{Z_4}(v)), \\ Z_4 \oplus (v + J_{Z_4}(v)), \\ (Z_1 + Z_4) \oplus (v + J_{Z_4}(v)) \end{array} \right\} \in \mathbb{R}^{3,4} \oplus \mathfrak{v}_{\min}. \quad (14)$$

We not only show the existence of a pair (C, A) for all the 9 cases (12), (13) and (14), but we also show that it form an affine space of a suitable dimension according to the initial conditions.

3.3. Relation between A and C in $[A, J_w] = J_{C(w)}$. We can express A in the form

$$A = \sum_{i < k} a_{i,k} J_{Z_i} J_{Z_k}. \quad (15)$$

Indeed, we know that A belongs to linear combination of $\{[J_{Z_i}, J_{Z_k}] : 1 \leq i < k \leq 7\}$, see details in [14, Proposition 4]. On the other hand, $[J_{Z_i}, J_{Z_k}] = J_{Z_i}J_{Z_k} - J_{Z_k}J_{Z_i} = 2J_{Z_i}J_{Z_k}$.

Then $[A, J_{Z_1}] = J_{C(Z_1)}$ implies

$$\begin{aligned} [A, J_{Z_1}] &= AJ_{Z_1} - J_{Z_1}A = \sum_{i < k} a_{i,k} J_{Z_i} J_{Z_k} J_{Z_1} - J_{Z_1} \sum_{i < k} a_{i,k} J_{Z_i} J_{Z_k} \\ &= 2 \sum_{k > 1} a_{1,k} J_{Z_k} = J_{2 \sum_{k=2}^7 a_{1,k} Z_k}. \end{aligned}$$

Thus $C(Z_1) = 2 \sum_{k=2}^7 a_{1,k} Z_k$. By similar calculations we obtain

$$\begin{aligned} C(Z_2) &= -2a_{1,2}Z_1 + 2 \sum_{k=3}^7 a_{2,k}Z_k, & C(Z_3) &= -2 \sum_{k=1}^2 a_{k,3}Z_k + 2 \sum_{k=4}^7 a_{3,k}Z_k, \\ C(Z_4) &= 2 \sum_{k=1}^3 a_{k,4}Z_k - 2 \sum_{k=5}^7 a_{4,k}Z_k, & C(Z_5) &= 2 \sum_{k=1}^4 a_{k,5}Z_k - 2 \sum_{k=5}^6 a_{5,k}Z_k, \\ C(Z_6) &= 2 \sum_{k=1}^5 a_{k,6}Z_k - 2a_{6,7}Z_7, & C(Z_7) &= 2 \sum_{k=1}^6 a_{k,7}Z_k. \end{aligned}$$

In the matrix form C is given by

$$C = 2 \times \begin{pmatrix} 0 & -a_{1,2} & -a_{1,3} & a_{1,4} & a_{1,5} & a_{1,6} & a_{1,7} \\ a_{1,2} & 0 & -a_{2,3} & a_{2,4} & a_{2,5} & a_{2,6} & a_{2,7} \\ a_{1,3} & a_{2,3} & 0 & a_{3,4} & a_{3,5} & a_{3,6} & a_{3,7} \\ a_{1,4} & a_{2,4} & a_{3,4} & 0 & a_{4,5} & a_{4,6} & a_{4,7} \\ a_{1,5} & a_{2,5} & a_{3,5} & -a_{4,5} & 0 & a_{5,6} & a_{5,7} \\ a_{1,6} & a_{2,6} & a_{3,6} & -a_{4,6} & -a_{5,6} & 0 & a_{6,7} \\ a_{1,7} & a_{2,7} & a_{3,7} & -a_{4,7} & -a_{5,7} & -a_{6,7} & 0. \end{pmatrix}. \quad (16)$$

3.4. Writing $A(v)$ and $A(J_{Z_4}(v))$ in the basis (8). Note that condition (7) implies

$$\begin{aligned} J_{Z_1}(v) &= -J_{Z_2}J_{Z_3}(v) = J_{Z_4}J_{Z_7}(v) = J_{Z_5}J_{Z_6}(v), \\ J_{Z_2}(v) &= J_{Z_1}J_{Z_3}(v) = J_{Z_4}J_{Z_6}(v) = -J_{Z_5}J_{Z_7}(v), \\ J_{Z_3}(v) &= -J_{Z_1}J_{Z_2}(v) = J_{Z_4}J_{Z_5}(v) = J_{Z_6}J_{Z_7}(v), \\ J_{Z_4}(v) &= J_{Z_1}J_{Z_7}(v) = J_{Z_2}J_{Z_6}(v) = J_{Z_3}J_{Z_5}(v), \\ J_{Z_5}(v) &= -J_{Z_2}J_{Z_7}(v) = -J_{Z_3}J_{Z_4}(v) = J_{Z_1}J_{Z_6}(v), \\ J_{Z_6}(v) &= -J_{Z_1}J_{Z_5}(v) = -J_{Z_2}J_{Z_4}(v) = J_{Z_3}J_{Z_7}(v), \\ J_{Z_7}(v) &= -J_{Z_1}J_{Z_4}(v) = J_{Z_2}J_{Z_5}(v) = -J_{Z_3}J_{Z_6}(v). \end{aligned} \quad (17)$$

Put $A(v) = \sum_{i < k} a_{i,k} J_{Z_i} J_{Z_k}(v) = \alpha_0 v + \sum_{j=1}^7 \alpha_j J_{Z_j}(v)$. Then using (17) we find the relations between coefficients $a_{i,k}$ and α_j as follows

$$\begin{cases} \alpha_0 = 0, \\ \alpha_1 = -a_{2,3} + a_{4,7} + a_{5,6}, \\ \alpha_2 = a_{1,3} + a_{4,6} - a_{5,7}, \\ \alpha_3 = -a_{1,2} + a_{4,5} + a_{6,7}, \\ \alpha_4 = a_{1,7} + a_{2,6} + a_{3,5}, \\ \alpha_5 = a_{1,6} - a_{2,7} - a_{3,4}, \\ \alpha_6 = -a_{1,5} - a_{2,4} + a_{3,7}, \\ \alpha_7 = -a_{1,4} + a_{2,5} - a_{3,6}. \end{cases} \quad (18)$$

Write $A(J_{Z_4}(v)) = \beta_0 v + \sum_{m=1}^7 \beta_m J_{Z_m}(v)$, and note that

$$\begin{aligned}
J_{Z_1}(v) &= J_{Z_1} J_{Z_4} J_{Z_4}(v) = J_{Z_2} J_{Z_5} J_{Z_4}(v) = -J_{Z_3} J_{Z_6} J_{Z_4}(v), \\
J_{Z_2}(v) &= -J_{Z_1} J_{Z_5} J_{Z_4}(v) = J_{Z_2} J_{Z_4} J_{Z_4}(v) = J_{Z_3} J_{Z_7} J_{Z_4}(v), \\
J_{Z_3}(v) &= J_{Z_1} J_{Z_6} J_{Z_4}(v) = -J_{Z_2} J_{Z_7} J_{Z_4}(v) = J_{Z_3} J_{Z_4} J_{Z_4}(v), \\
v &= J_{Z_1} J_{Z_7} J_{Z_4}(v) = J_{Z_2} J_{Z_6} J_{Z_4}(v) = J_{Z_3} J_{Z_5} J_{Z_4}(v), \\
J_{Z_5}(v) &= -J_{Z_1} J_{Z_2} J_{Z_4}(v) = -J_{Z_4} J_{Z_5} J_{Z_4}(v) = J_{Z_6} J_{Z_7} J_{Z_4}(v), \\
J_{Z_6}(v) &= J_{Z_1} J_{Z_3} J_{Z_4}(v) = -J_{Z_4} J_{Z_6} J_{Z_4}(v) = -J_{Z_5} J_{Z_7} J_{Z_4}(v), \\
J_{Z_7}(v) &= -J_{Z_2} J_{Z_3} J_{Z_4}(v) = -J_{Z_4} J_{Z_7} J_{Z_4}(v) = J_{Z_5} J_{Z_6} J_{Z_4}(v).
\end{aligned} \tag{19}$$

By using (19) we find relations between $a_{i,k}$ and β_j as below:

$$\begin{cases}
\beta_0 = a_{1,7} + a_{2,6} + a_{3,5}, \\
\beta_1 = a_{1,4} + a_{2,5} - a_{3,6}, \\
\beta_2 = -a_{1,5} + a_{2,4} + a_{3,7}, \\
\beta_3 = a_{1,6} - a_{2,7} + a_{3,4}, \\
\beta_4 = 0, \\
\beta_5 = -a_{1,2} - a_{4,5} + a_{6,7}, \\
\beta_6 = a_{1,3} - a_{4,6} - a_{5,7}, \\
\beta_7 = -a_{2,3} - a_{4,7} + a_{5,6}.
\end{cases} \tag{20}$$

3.5. Existence of $D = (C, A)$. Based on data in the previous section, we discuss the existence of a pair (C, A) of condition (9) for each cases of initial conditions (12), (13), and (14).

CASE (z_0, x_0) WITH $\|x_0\|^2 \neq 0$. Assume for the moment that $\|x_0\|^2 = \|z_0\|^2 = 1$ and use Lemma 1. Then the conditions $A(v) = J_{Z_1}(v)$, $C(Z_1) = 0$, and (18) give

$$a_{1,2} = a_{1,3} = a_{1,4} = a_{1,5} = a_{1,6} = a_{1,7} = 0, \quad \text{and}$$

$$\begin{cases}
\alpha_0 = 0, \\
\alpha_1 = -a_{2,3} + a_{4,7} + a_{5,6} = 1, \\
\alpha_2 = a_{1,3} + a_{4,6} - a_{5,7} = 0, \\
\alpha_3 = -a_{1,2} + a_{4,5} + a_{6,7} = 0, \\
\alpha_4 = a_{1,7} + a_{2,6} + a_{3,5} = 0, \\
\alpha_5 = a_{1,6} - a_{2,7} - a_{3,4} = 0, \\
\alpha_6 = -a_{1,5} - a_{2,4} + a_{3,7} = 0, \\
\alpha_7 = -a_{1,4} + a_{2,5} - a_{3,6} = 0.
\end{cases} \implies \begin{cases}
a_{5,6} = 1 + a_{2,3} - a_{4,7}, \\
a_{5,7} = a_{4,6}, \\
a_{6,7} = -a_{4,5}, \\
a_{3,5} = -a_{2,6}, \\
a_{3,4} = -a_{2,7}, \\
a_{3,7} = a_{2,4}, \\
a_{3,6} = a_{2,5},
\end{cases} \tag{21}$$

where $a_{2,3}, a_{2,4}, a_{2,5}, a_{2,6}, a_{2,7}, a_{4,5}, a_{4,6}, a_{4,7}$ are free parameters.

Analogously to the above case one can obtain systems similar to (21) for any of the following initial conditions:

$$Z_1 \oplus v, \quad Z_4 \oplus v, \quad Z_1 \oplus J_{Z_4}(v), \quad Z_4 \oplus J_{Z_4}(v), \quad (Z_1 + Z_4) \oplus v, \quad (Z_1 + Z_4) \oplus J_{Z_4}(v). \tag{22}$$

CASE (z_0, x_0) WITH $\|x_0\| = 0$ OR $\|z_0\| = 0$. The cases $Z_1 \oplus (v + J_{Z_4}(v))$, $Z_4 \oplus (v + J_{Z_4}(v))$, and $(Z_1 + Z_4) \oplus (v + J_{Z_4}(v))$ are treated by a similar way, requiring more careful calculation. We consider only the $(Z_1 + Z_4) \oplus (v + J_{Z_4}(v))$ in some details.

Conditions (9) are equivalent to

$$\begin{aligned} A(v + J_{Z_4}(v)) &= J_{Z_1+Z_4}(v + J_{Z_4}(v)) \\ &= J_{Z_1}(v) + J_{Z_1Z_4}(v) + J_{Z_4}(v) + v = \gamma_0 v + \sum_{i=1}^7 \gamma_i J_{Z_i}(v). \end{aligned}$$

Noticing that $J_{Z_1}(v) + J_{Z_1Z_4}(v) + J_{Z_4}(v) + v = J_{Z_1}(v) - J_{Z_7}(v) + J_{Z_4}(v) + v$, we obtain the system

$$\begin{cases} \gamma_0 = \alpha_0 + \beta_0 = a_{1,7} + a_{2,6} + a_{3,5} = 1, \\ \gamma_1 = \alpha_1 + \beta_1 = (-a_{2,3} + a_{5,6} - a_{3,6} + a_{2,5}) + (a_{1,4} + a_{4,7}) = 1, \\ \gamma_2 = \alpha_2 + \beta_2 = (a_{1,3} - a_{1,5} + a_{3,7} - a_{5,7}) + (a_{4,6} + a_{2,4}) = 0, \\ \gamma_3 = \alpha_3 + \beta_3 = (-a_{1,2} + a_{6,7} + a_{1,6} - a_{2,7}) + (a_{4,5} + a_{3,4}) = 0, \\ \gamma_4 = \alpha_4 + \beta_4 = a_{1,7} + a_{2,6} + a_{3,5} = 1, \\ \gamma_5 = \alpha_5 + \beta_5 = (-a_{1,2} + a_{6,7} + a_{1,6} - a_{2,7}) - (a_{4,5} + a_{3,4}) = 0, \\ \gamma_6 = \alpha_6 + \beta_6 = (a_{1,3} - a_{1,5} + a_{3,7} - a_{5,7}) - (a_{4,6} + a_{2,4}) = 0, \\ \gamma_7 = \alpha_7 + \beta_7 = (-a_{2,3} + a_{5,6} - a_{3,6} + a_{2,5}) - (a_{1,4} + a_{4,7}) = -1. \end{cases} \quad (23)$$

The condition $C(Z_1 + Z_4) = 0$ leads to

$$a_{1,4} = 0, \quad a_{1,2} + a_{2,4} = 0, \quad a_{1,3} + a_{3,4} = 0, \quad a_{1,5} - a_{4,5} = 0, \quad a_{1,6} - a_{4,6} = 0, \quad a_{1,7} - a_{4,7} = 0,$$

From these expressions we obtain the solution

$$\begin{cases} a_{1,4} = 0, \\ a_{4,6} = -a_{2,4} = a_{1,6} = a_{1,2}, \\ a_{4,5} = -a_{3,4} = a_{1,5} = a_{1,3}, \\ a_{4,7} = a_{1,7} = 1, \\ a_{5,6} = a_{2,3} - a_{2,5} + a_{3,6}, \\ a_{3,5} = -a_{2,6}, \\ a_{6,7} = a_{2,7}, \\ a_{5,7} = a_{3,7}, \end{cases}$$

with eight free parameters $a_{1,2}, a_{1,3}, a_{2,3}, a_{2,5}, a_{2,6}, a_{2,7}, a_{3,6}, a_{3,7}$.

This finishes the proof of Theorem 4. We emphasize that this proof is constructive and it is easy to find a 1-parameter group (and even a multi-parameter family of such groups) whose orbit is an arbitrarily chosen geodesic.

4. Riemannian H -type Nilmanifold $N_{7,0} = N_{7,0}(\mathfrak{v}_{\min})$

Here we consider another example of 15-dimensional manifold, related to Clifford algebra. It is the Riemannian H -type Lie group with 7-dimensional centre and 8-dimensional irreducible Clifford module. The metric on the centre and on the module is positive definite. It is known from the work of [13] that this Riemannian nilmanifold $N_{7,0}(\mathfrak{v}_{\min})$ is geodesic orbit. We give a proof which is similar to the proof in Section 3, but significantly simpler

since all the vectors has positive square lengths. It also shows more complicated structure of pseudo-Riemannian nilmanifold, compared to Riemannian once.

Theorem 5. *The Riemannian H -type nilmanifold $N_{7,0}(\mathfrak{v}_{\min})$ is geodesic orbit. Each geodesic is realized as an orbit of a one-parameter subgroup which initial conditions form an eight dimensional affine space.*

◁ The Clifford algebra $\text{Cl}(\mathbb{R}^{7,0})$ has two non-equivalent irreducible 8-dimensional modules \mathfrak{v}_{\min}^+ and \mathfrak{v}_{\min}^- . The corresponding 15-dimensional H -type Lie algebras $N_{7,0}(\mathfrak{v}_{\min}^+)$ and $N_{7,0}(\mathfrak{v}_{\min}^-)$ are isomorphic and isometric, see [25, Section 3.1.2]. The left invariant metric is generated by the positive definite scalar product $\langle \cdot, \cdot \rangle_{7,0} + \langle \cdot, \cdot \rangle_{8,0}$ on the Lie algebra $\mathfrak{n}_{7,0} \cong \mathbb{R}^{7,0} \oplus \mathbb{R}^{8,0}$. We write $(z_0, x_0) \in \mathfrak{z} \oplus \mathfrak{v}_{\min} \cong \mathbb{R}^{7,0} \oplus \mathbb{R}^{8,0}$ for an initial condition for a geodesic. As was shown in Section 3, without loss of generality we can assume that $z_0 = Z_1$, where

$$\{Z_1, \dots, Z_7\}, \quad \|Z_1\|_{\mathbb{R}^{7,0}}^2 = \dots = \|Z_7\|_{\mathbb{R}^{7,0}}^2 = 1,$$

is an orthonormal basis for $\mathbb{R}^{7,0}$ and $x_0 = v \in \mathfrak{v}_{\min}$ is a vector satisfying (7) and $\|v\|_{\mathbb{R}^{8,0}}^2 = 1$. We will also use basis (8).

Then, a geodesic determined by the initial condition $(z_0, x_0) = (Z_1, v)$ is homogeneous, if there is a pair $D = (C, A)$, such that

$$A(v) = J_{Z_1}(v), \quad C(Z_1) = 0, \quad [A, J_w] = J_{C(w)} \quad \text{for all } w \in \mathbb{R}^{7,0}. \quad (24)$$

We write $A = \sum_{i < k} a_{i,k} J_{Z_i} J_{Z_k}$. By checking the condition $[A, J_w] = J_{C(w)}$ for $w = Z_k$, $k = 1, \dots, 7$, we come to the conclusion that

$$C = 2 \times \begin{pmatrix} 0 & -a_{1,2} & -a_{1,3} & -a_{1,4} & -a_{1,5} & -a_{1,6} & -a_{1,7} \\ a_{1,2} & 0 & -a_{2,3} & -a_{2,4} & -a_{2,5} & -a_{2,6} & -a_{2,7} \\ a_{1,3} & a_{2,3} & 0 & -a_{3,4} & -a_{3,5} & -a_{3,6} & -a_{3,7} \\ a_{1,4} & a_{2,4} & a_{3,4} & 0 & -a_{4,5} & -a_{4,6} & -a_{4,7} \\ a_{1,5} & a_{2,5} & a_{3,5} & a_{4,5} & 0 & -a_{5,6} & -a_{5,7} \\ a_{1,6} & a_{2,6} & a_{3,6} & a_{4,6} & a_{5,6} & 0 & -a_{6,7} \\ a_{1,7} & a_{2,7} & a_{3,7} & a_{4,7} & a_{5,7} & a_{6,7} & 0 \end{pmatrix}. \quad (25)$$

Note that condition (7) for $N_{7,0}(\mathfrak{v}_{\min})$ implies

$$\begin{aligned} J_{Z_1}(v) &= -J_{Z_2} J_{Z_3}(v) = -J_{Z_4} J_{Z_7}(v) = -J_{Z_5} J_{Z_6}(v), \\ J_{Z_2}(v) &= J_{Z_1} J_{Z_3}(v) = J_{Z_4} J_{Z_6}(v) = -J_{Z_5} J_{Z_7}(v), \\ J_{Z_3}(v) &= -J_{Z_1} J_{Z_2}(v) = J_{Z_4} J_{Z_5}(v) = J_{Z_6} J_{Z_7}(v), \\ J_{Z_4}(v) &= J_{Z_1} J_{Z_7}(v) = -J_{Z_2} J_{Z_6}(v) = -J_{Z_3} J_{Z_5}(v), \\ J_{Z_5}(v) &= J_{Z_2} J_{Z_7}(v) = J_{Z_3} J_{Z_4}(v) = J_{Z_1} J_{Z_6}(v), \\ J_{Z_6}(v) &= -J_{Z_1} J_{Z_5}(v) = J_{Z_2} J_{Z_4}(v) = -J_{Z_3} J_{Z_7}(v), \\ J_{Z_7}(v) &= -J_{Z_1} J_{Z_4}(v) = -J_{Z_2} J_{Z_5}(v) = J_{Z_3} J_{Z_6}(v). \end{aligned} \quad (26)$$

Now we write the value $A(v)$ in the basis (8). Put $A(v) = \sum_{i < k} a_{i,k} J_{Z_i} J_{Z_k}(v) = \alpha_0 v + \sum_{j=1}^7 \alpha_j J_{Z_j}(v)$ and solve the equation $A(v) = J_{Z_1}(v)$. We obtain the values of α_k in terms of $a_{j,k}$.

$$\begin{cases} \alpha_0 = 0, \\ \alpha_1 = -a_{2,3} - a_{4,7} - a_{5,6} = 1, \\ \alpha_2 = a_{1,3} + a_{4,6} - a_{5,7} = 0, \\ \alpha_3 = -a_{1,2} + a_{4,5} + a_{6,7} = 0, \\ \alpha_4 = a_{1,7} - a_{2,6} - a_{3,5} = 0, \\ \alpha_5 = a_{1,6} + a_{2,7} + a_{3,4} = 0, \\ \alpha_6 = -a_{1,5} + a_{2,4} - a_{3,7} = 0, \\ \alpha_7 = -a_{1,4} - a_{2,5} + a_{3,6} = 0. \end{cases} \quad (27)$$

The condition $C(Z_1) = 0$ implies

$$a_{1,2} = a_{1,3} = a_{1,4} = a_{1,5} = a_{1,6} = a_{1,7} = 0.$$

The system (27) has the following solution:

$a_{2,3}, a_{2,4}, a_{2,5}, a_{2,6}, a_{2,7}, a_{4,5}, a_{4,6}, a_{4,7}$ are free variables and

$$\begin{cases} a_{3,4} = -a_{2,7}, \\ a_{3,5} = -a_{2,6}, \\ a_{3,6} = a_{2,5}, \\ a_{3,7} = a_{2,4}, \\ a_{5,6} = 1 + a_{2,3} - a_{4,7}, \\ a_{5,7} = a_{4,6}, \\ a_{6,7} = -a_{4,5}. \end{cases}$$

This finishes the proof of Theorem 5. \triangleright

The given examples of geodesic orbit pseudo-Riemannian metrics are obtained using important structural results for the corresponding pseudo H -type Lie algebras. We dare to suggest that similar ideas will be useful in constructing other examples of geodesic orbit pseudo-Riemannian manifolds.

References

1. Kowalski, O. and Vanhecke, L. Riemannian Manifolds with Homogeneous Geodesics, *Bollettino dell'Unione Matematica Italiana. B* (7), 1991, vol. 5, no. 1, pp. 189–246.
2. Arvanitoyeorgos, A. Homogeneous Manifolds Whose Geodesics are Orbits. Recent Results and Some Open Problems, *Irish Mathematical Society Bulletin*, 2017, vol. 79, pp. 5–29. DOI: 10.33232/BIMS.0079.5.29.
3. Berestovskii, V. N. and Nikonorov, Yu. G. *Riemannian Manifolds and Homogeneous Geodesics*, Springer Monographs in Mathematics, Cham, Springer, 2020.
4. Nikolayevsky, Y. and Ziller, W. Non-Singular Geodesic Orbit Nilmanifolds, *Annali di Matematica Pura ed Applicata*, 2026. DOI: 10.1007/s10231-026-01661-9.
5. Nikonorov, Yu. G. On the Structure of Geodesic Orbit Riemannian Spaces, *Annals of Global Analysis and Geometry*, 2017, vol. 52, pp. 289–311. DOI: 10.1007/s10455-017-9558-0.
6. Nikonorov, Yu. G. On Geodesic Orbit Nilmanifolds, *Journal of Geometry and Physics*, 2024, vol. 203, article ID 105257, 12 p.
7. del Barco, V. Homogeneous Geodesics in Pseudo-Riemannian Nilmanifolds, *Advances in Geometry*, 2016, vol. 16, no. 2, pp. 175–187.
8. Chen, Z., Nikolayevsky, Y., Wolf, J. A. and Zhang, S. Pseudo-Riemannian Geodesic Orbit Nilmanifolds of Signature $(n - 2, 2)$, *The Journal of Geometric Analysis*, 2024, vol. 34, no. 5, article no. 132, 21 p. DOI: 10.1007/s12220-024-01579-9.
9. Chen, Z., Wolf, J. A. and Zhang, S. On the Geodesic Orbit Property for Lorentz Manifolds, *The Journal of Geometric Analysis*, 2022, vol. 32, no. 3, article no. 81, 14 p. DOI: 10.1007/s12220-021-00744-8.

10. Dušek, Z. and Kowalski, O. Light-like Homogeneous Geodesics and the Geodesic Lemma for Any Signature, *Publicationes Mathematicae Debrecen*, 2007, vol. 71, no. 1–2, pp. 245–252. DOI: 10.5486/PMD.2007.3800.
11. Nikolayevsky, Y. and Wolf, J. A. The Structure of Geodesic Orbit Lorentz Nilmanifolds, *The Journal of Geometric Analysis*, 2023, vol. 33, no. 3, article no. 82, 12 p. DOI: 10.1007/s12220-022-01134-4.
12. Kaplan, A. On the Geometry of Groups of Heisenberg Type, *Bulletin of the London Mathematical Society*, 1983, vol. 15, pp. 35–42. DOI: 10.1112/blms/15.1.35.
13. Riehm, C. Explicit Spin Representations and Lie Algebras of Heisenberg Type, *Journal of the London Mathematical Society*, 1984, vol. 29, no. 1, pp. 49–62. DOI: 10.1112/jlms/s2-29.1.49.
14. Furutani, K., Markina, I. and Nikonorov, Y. *Geodesic Orbit Pseudo-Riemannian H-Type Nilmanifolds: Case of Minimal Admissible Clifford Modules*, Preprint, 2025, arXiv: 2507.00470.
15. Lee, J. M. *Introduction to Riemannian Manifolds*, 2nd edition, Graduate Texts in Mathematics, vol. 176, Cham, Springer, 2018.
16. Cordero, L. A. and Parker, P. E. Isometry Groups of Pseudoriemannian 2-Step Nilpotent Lie Groups, *Houston Journal of Mathematics*, 2009, vol. 35, no. 1, pp. 49–72.
17. Gordon, C. Homogeneous Riemannian Manifolds Whose Geodesics are Orbits, *Topics in Geometry: Honoring the Memory of Joseph D’Atri*, Progress in Nonlinear Differential Equations, vol. 20, Birkhäuser, 1996, pp. 155–174.
18. Ovando, G. P. Naturally Reductive Pseudo-Riemannian 2-Step Nilpotent Lie Groups, *Houston Journal of Mathematics*, 2013, vol. 39, no. 1, pp. 147–167.
19. Kaplan, A. Fundamental Solutions for a Class of Hypoelliptic PDE Generated by Composition of Quadratic Forms, *Transactions of the American Mathematical Society*, 1980, vol. 258, no. 1, pp. 147–153. DOI: 10.1090/S0002-9947-1980-0554324-X.
20. Ciatti, P. Scalar Products on Clifford Modules and Pseudo- H -Type Lie Algebras, *Annali di Matematica Pura ed Applicata*, IV Ser., 2000, vol. 178, pp. 1–31. DOI: 10.1007/BF02505885.
21. Godoy Molina, M., Korolko, A. and Markina, I. Sub-Semi-Riemannian Geometry of General H -Type Groups, *Bulletin des Sciences Mathématiques*, 2013, vol. 137, no. 6, pp. 805–833. DOI: 10.1016/j.bulsci.2013.05.005.
22. Kaplan, A. Riemannian Nilmanifolds Attached to Clifford Modules, *Geometriae Dedicata*, 1981, vol. 11, pp. 127–136. DOI: 10.1007/BF00147615.
23. Furutani, K. and Markina, I. Automorphism Groups of Pseudo H -Type Algebras, *Journal of Algebra*, 2021, vol. 568, no. 3, pp. 91–138. DOI: 10.1016/j.jalgebra.2020.09.038.
24. Furutani, K. and Markina, I. Complete Classification of Pseudo H -type Lie Algebras. I, *Geometriae Dedicata*, 2017, vol. 190, pp. 23–51. DOI: 10.1007/s10711-017-0225-1.
25. Berndt, J., Tricerri, F. and Vanhecke, L. *Generalized Heisenberg Groups and Damek–Ricci Harmonic Spaces*, Lecture Notes in Mathematics, vol. 1598, Berlin, Springer, 1995.

Received November 8, 2025

IRINA G. MARKINA
 Department of Mathematics, University of Bergen,
 P. O. Box 7803, N-5020 Bergen, 41 Allégaten, Norway,
 Professor
 E-mail: irina.markina@uib.no
<https://orcid.org/0000-0003-2686-6657>

YURIĬ G. NIKONOROV
 Southern Mathematical Institute of VSC RAS,
 53 Vatutina St., Vladikavkaz 362025, Russia,
 Principal Investigator
 E-mail: nikonorov2006@mail.ru
<https://orcid.org/0000-0002-9671-2314>

KENRO FURUTANI
 Osaka Central Advanced Mathematical Institute,
 Osaka Metropolitan University,
 Sugimoto, Sumiyoshi-ku, Osaka 558-8585, Japan,
 Professor
 E-mail: kf46089@gmail.com
<https://orcid.org/0000-0002-9972-7693>

О ПРИМЕРАХ ПСЕВДОРИМАНОВЫХ ГЕОДЕЗИЧЕСКИ
ОРБИТАЛЬНЫХ МНОГООБРАЗИЙМаркина И. Г.¹, Никоноров Ю. Г.², Фурутани К.³¹ Университет Бергена, Норвегия, 5020, Берген, Аллегатен, 41;² Южный математический институт — филиал ВНИИ РАН,
Россия, 362025, Владикавказ, ул. Ватутина, 53;³ Столичный университет Осаки,
Осакский центральный институт углубленной математики,
Япония, 558-8585, Осака, Сугимото, Сумиоши-ку

E-mail: irina.markina@uib.no, nikonorov2006@mail.ru, kf46089@gmail.com

Аннотация. Псевдориманово многообразие (M, g) называется геодезически орбитальным многообразием, если любая геодезическая γ многообразия M является орбитой 1-параметрической подгруппы полной группы изометрий (M, g) . Этот термин в случае римановых многообразий был введен в 1991 г. О. Ковальским и Л. Ванхекке, положившими начало систематическому изучению пространств $(M = G/H, g)$, где G — группа изометрий, а H — подгруппа изотропии. Следует отметить, что симметричные пространства, слабо симметричные пространства, естественно редуктивные однородные пространства, нормальные однородные пространства, обобщенные нормальные однородные пространства (и не только они) являются подклассами класса геодезически орбитальных псевдоримановых пространств. В данной статье приводятся примеры псевдоримановых геодезически орбитальных многообразий. Таковыми примерами являются специальные 15-мерные группы Ли псевдо H -типа, т. е. 2-ступенно нильпотентные группы Ли гейзенберговского типа, снабженные левоинвариантной псевдоримановой метрикой. Для построения соответствующих примеров использованы результаты о структуре рассматриваемых групп Ли.

Ключевые слова: римановы геодезически орбитальные многообразия, псевдоримановы геодезически орбитальные многообразия, группы Ли типа Гейзенберга.

AMS Subject Classification: 53C50, 53C30, 22E25.

Образец цитирования: Markina I. G., Nikonorov Yu. G. and Furutani K. On Examples of Geodesic Orbit Pseudo-Riemannian Manifolds // Владикавк. мат. журн.—2026.—Т. 28, № 1.—С. 108–121 (in English). DOI: 10.46698/i4125-5722-6924-j.